



## **DRY MOUTH** (xerostomia or hyposialia)

### **Mouth spray**

### **Lubricating and protective solution for intraoral sprays.**

#### **Presentation:**

Aluminium bottle with 0.1 ml dosing pump containing 40 ml of solution, i.e. 400 doses.

#### **Composition:**

Triesters of Glycerol Oxidized of essential fatty acids of vegetable origin (O.G.T.): 94.4%  
Silicon dioxide  
Food-grade flavouring: orange-grapefruit and mint  
Aspartame

#### **What is xerostomia?**

Xerostomia is excessive dryness of the mouth. It is caused by secretion of saliva that is insufficient (hyposialia) or even nonexistent (asialia). It may be temporary or prolonged, depending on the cause: anxiety, dehydration, medicines, radiotherapy, systemic illnesses. It requires medical monitoring.

#### **The role of aequasyl**

Xerostomia has damaging effects on the oral cavity and on quality of life. For this reason, treatment of dryness of the mouth is based primarily on relieving the associated symptoms.

Controlled clinical trials demonstrate that aequasyl quickly and significantly improves the symptoms accompanying dryness of the mouth. The effects of aequasyl are:

- lubricating and adhesive, creating a lipid film designed to limit the loss of water and restore the viscoelasticity of the oral mucosa,
- protecting against local aggressions.

#### **Indication:**

In adults, aequasyl improves the symptoms accompanying xerostomia or hyposialia (dry mouth), in particular where this is caused by medicinal products or radiotherapy.

#### **Instructions for use:**

Shake gently before use  
During first use: prime the pump by pressing the push-button.  
One intraoral spray on the inside of each cheek three to four times daily.

Following each spray, gently move the product to inflamed areas using the tongue.

#### **Precautions for use:**

This product has been shown to be non-toxic and well tolerated. It is not a drug product, it is a medical device.

If you use aequasyl, you should also take any other treatments prescribed by your doctor and your condition should be monitored by medical personnel.

If you experience an allergic reaction, it is recommended that you discontinue treatment as a precautionary measure and consult your doctor.

For adults only.

External use, intraoral spray.

#### **Keep out of the reach of children.**

Do not apply to the eyes.

This product does not contain any, propellant gas.

Contains aspartame.

#### **Storage conditions:**

Store at room temperature.

Keep the bottle upright, if stored horizontally the fluid may leak.

#### **Manufacturer:**

Laboratoires Carilene, 7 rue du Chant des Oiseaux, F-78360 Montesson, FRANCE

#### **Distributor:**

**bluechip**  
HEALTHCARE

Bluechip Healthcare Marketing Ltd,  
34 South Molton St, London, W1K 5RG UK.

Medical Device

CE marking: 2005

40ml

